

Certainty Quackgrass Trial 2006

Nick Christians, Christopher J. Blume, and Ryan Krull

Objectives: The objectives of this study were to observe the effects of Certainty 75 WG (sulfosulfuron), a Monsanto product, for the control of quackgrass (*Elymus repens*) in Kentucky bluegrass turf.

Methods: A mature stand of South Dakota Common Kentucky bluegrass with a uniform infestation of quackgrass was chosen for the study. Plots measured 5 ft x 5 ft. A single application of certainty at levels from 0.0234 to 0.47 lb. ai/acre with a nonionic surfactant (X-77) at 0.25% v/v, were applied on July 6, 2006. Both treatments 2 and 3 were applied at the same level of 0.0234 lb. ai/acre. No phytotoxicity was observed on the Kentucky bluegrass at any time following treatment. Data were collected on % cover of quackgrass at the end of the season on Oct. 24 with a grid. The 3 x 3 ft grid had strings arranged so that there were 100 locations where the strings crossed. Wherever quackgrass occurred below the crossed strings it was counted. Quackgrass was reported on a percentage cover of the plot.

The Certainty reduced quackgrass in all treated plots as compared to the control at the end of the season by 75 to 92 % (Table 1).

Treatment	Rate of Certainty lb a.i./acre	Rate/plot	Phytotoxicity 7/14	Quackgrass % Cover
1. Control		-	9	12
2. Certainty	0.0234*	.008	9	3
3. Certainty	0.0234	.008	9	3
4. Certainty	0.035	.012	9	1
5. Certainty	0.047	.016	9	2
LSD 0.05			NS	4

Phytotoxicity based on a scale of 9 to 1, where 9=no damage.
* each treatment was applied with 0.25 % nonionic surfactant (v/v) (X-77)