



R= 137634

### 1999 Corn Gluten Meal/Urea Crabgrass Control Study - Year 9-2007

Nick Christians, Christopher J. Blume, and Matt Klingenberg

This study was initiated in 1999 to determine if the levels of annual grass and broadleaf weed control provided by corn gluten meal (CGM) treatments can be explained by the nitrogen response of treated bluegrass and not herbicidal activity of CGM. The study is being conducted at the Iowa State University Research Station north of Ames, IA in an area of 'Parade' Kentucky bluegrass. The soil in this experimental area is a Nicollet (fine-loamy, mixed, mesic Aquic Hapludolls) with an organic matter content of 4.2%, a pH of 6.75, 17 ppm P, and 103 ppm K.

The experimental design is a randomized complete block with three replications. Individual experimental plots are 5 x 5 ft with five treatments. Corn gluten meal and urea are applied yearly to the same plots at an annual rate of 4 lbs N/1000 ft<sup>2</sup> (Table 1). Treatments include split applications of 2 lb N/1000 ft<sup>2</sup> and four applications of 1 lb N/1000 ft<sup>2</sup> each. The CGM and urea are applied using cardboard containers as 'shaker dispensers'. The materials are watered-in with the irrigation system. Supplemental irrigation is used to provide adequate moisture to maintain the grass in good growing condition. In 2007, initial applications of all urea and CGM treatments were made on April 26. Sequential applications of 1 lb N/1000 ft<sup>2</sup> were made on June 11, August 6, and September 20. The second applications of 2 lb N/1000 ft<sup>2</sup> for urea and CGM (Treatment 3 and 5) were made on August 6.

Turf quality was monitored from May through September (Table 1). Visual turf quality was assessed using a 9 to 1 scale with 9 = best, 6 = lowest acceptable, and 1 = worst turf quality.

Crabgrass data represent the percentage of plants per individual plot. Crabgrass assessments were made on July 31 and August 31 (Table 2).

Broadleaf weed populations were measured by either counting the number of plants or estimating the percentage cover per individual plot. Data for dandelion and clover were taken beginning in May and ending in September. Dandelion infestations were determined by counting the number of plants per individual plot. Clover populations were estimated by assessing the percentage area of each plot covered by clover.

Data were analyzed with the Statistical Analysis System (SAS) and the General Linear Model (GLM) procedure. Effects of CGM and urea on bluegrass quality and weed control were examined using Fisher's Least Significant Difference (LSD) means comparison tests.

Table 1. Visual quality<sup>1</sup> of Kentucky bluegrass treated in the 1991 Corn Gluten Meal Weed Control Study.

Material	lbs N /1000 ft <sup>2</sup>	Number of applications	May 7	June 4	June 30	July 31	August 31	October 3
1 Untreated control	NA	NA	6.7	5.3	5.3	4.7	5.3	6.0
2 Corn gluten meal	4	4	6.7	5.7	5.7	6.0	6.0	6.3
3 Corn gluten meal	4	2	6.7	6.7	6.0	6.0	5.7	6.0
4 Urea (46-0-0)	4	4	5.7	5.0	6.0	5.0	4.3	5.7
5 Urea (46-0-0)	4	2	6.7	6.0	6.0	5.7	6.0	6.3
LSD <sub>0.05</sub>			NS	NS	NS	NS	NS	NS

<sup>1</sup>Visual quality was assessed using a 9 to 1 scale with 9 = best, 6 = lowest acceptable, and 1 = worst turf quality. NS = means are not significantly different at the 0.05 level.

Table 2. Crabgrass counts<sup>1</sup> in Kentucky bluegrass treated in the 1999 Corn Gluten Meal/Urea Weed Control Study.

Material	lbs N /1000 ft <sup>2</sup>	Number of applications	July 31	August 31
1 Untreated control	NA	NA	80	71.7
2 Corn gluten meal	4	4	46.7	31.7
3 Corn gluten meal	4	2	48.3	47.7
4 Urea (46-0-0)	4	4	53.3	53.3
5 Urea (46-0-0)	4	2	46.7	50.0
LSD <sub>0.05</sub>			NS	NS

<sup>1</sup>Values represent the number of crabgrass plants per plot covered. NS = means are not significantly different at the 0.05 level.

**Table 3.** Dandelion counts<sup>1</sup> in Kentucky bluegrass treated in the 1999 Corn Gluten Meal/Urea Weed Control Study.

	Material	lbs N /1000 ft <sup>2</sup>	Number of applications	May 7	June 4	June 30	July 31	August 31	October 3
1	Untreated control	NA	NA	18.7	15.7	19.7	12.0	24.0	25.0
2	Corn gluten meal	4	4	19.7	17.7	23.7	12.7	20.0	19.0
3	Corn gluten meal	4	2	16.0	12.7	15.3	11.7	19.7	20.7
4	Urea (46-0-0)	4	4	13.3	11.0	12.0	7.7	10.0	9.3
5	Urea (46-0-0)	4	2	10.0	7.7	10.3	3.7	8.0	10.0
<b>LSD<sub>0.05</sub></b>				NS	NS	14.4	NS	NS	NS

<sup>1</sup>Values represent the number of dandelion plants per plot.

NS = means are not significantly different at the 0.05 level.

**Table 4.** Percentage clover cover<sup>1</sup> in Kentucky bluegrass treated in the 1999 Corn Gluten Meal/Urea Weed Control Study.

	Material	lbs N /1000 ft <sup>2</sup>	Number of applications	May 7	June 4	June 30	July 31	August 31	October 3
1	Untreated control	NA	NA	18.3	5.0	4.0	11.7	3.7	2.7
2	Corn gluten meal	4	4	26.7	8.7	7.7	23.7	5.3	3.0
3	Corn gluten meal	4	2	15.0	7.7	5.7	16.0	3.3	1.3
4	Urea (46-0-0)	4	4	35.0	18.3	21.0	26.7	8.3	5.7
5	Urea (46-0-0)	4	2	25.0	18.3	21.3	43.3	8.0	3.7
<b>LSD<sub>0.05</sub></b>				NS	NS	NS	NS	NS	NS

<sup>1</sup>Values represent the area per plot covered by clover.

NS = means are not significantly different at the 0.05 level.